

Extremal Black Holes from Homotopy Algebras

Chettha Saelim

University of Surrey

Based on arXiv:2508.08886 with Jan Gutowski and Martin Wolf

- **BH uniqueness theorem** [Robinson; Mazur; Carter; Israel; Amsel, Horowitz, Marolf, and Roberts]: In 4D, a stationary asymptotically flat black hole is Kerr-Newman.
- The theorem breaks down in higher dimensions, e.g. in 5D, Myers–Perry solutions [Myers and Perry] and black ring solutions [Empanan and Reall] have the same set of conserved charges.
- **Deformation of near-horizon geometry** [Li and Lucietti; Katona and Lucietti; Fontanella and Gutowski]: Finiteness of linear transverse deformations. A black hole with a static and maximally symmetric near-horizon geometry is Schwarzschild–de Sitter.
- L_∞ -**algebra** (review [Jurčo, Raspollini, Sämann, and Wolf]) is a powerful framework in classical and quantum field theory. The **homological perturbation lemma** allows one to solve deformation problems systematically.

- **BH uniqueness theorem** [Robinson; Mazur; Carter; Israel; Amsel, Horowitz, Marolf, and Roberts]: In 4D, a stationary asymptotically flat black hole is Kerr-Newman.
- The theorem breaks down in higher dimensions, e.g. in 5D, Myers–Perry solutions [Myers and Perry] and black ring solutions [Empanan and Reall] have the same set of conserved charges.
- **Deformation of near-horizon geometry** [Li and Lucietti; Katona and Lucietti; Fontanella and Gutowski]: Finiteness of linear transverse deformations. A black hole with a static and maximally symmetric near-horizon geometry is Schwarzschild–de Sitter.
- L_∞ -**algebra** (review [Jurčo, Raspollini, Sämann, and Wolf]) is a powerful framework in classical and quantum field theory. The **homological perturbation lemma** allows one to solve deformation problems systematically.

- **BH uniqueness theorem** [Robinson; Mazur; Carter; Israel; Amsel, Horowitz, Marolf, and Roberts]: In 4D, a stationary asymptotically flat black hole is Kerr-Newman.
- The theorem breaks down in higher dimensions, e.g. in 5D, Myers–Perry solutions [Myers and Perry] and black ring solutions [Empanan and Reall] have the same set of conserved charges.
- **Deformation of near-horizon geometry** [Li and Lucietti; Katona and Lucietti; Fontanella and Gutowski]: Finiteness of linear transverse deformations. A black hole with a static and maximally symmetric near-horizon geometry is Schwarzschild–de Sitter.
- L_∞ -**algebra** (review [Jurčo, Raspollini, Sämann, and Wolf]) is a powerful framework in classical and quantum field theory. The **homological perturbation lemma** allows one to solve deformation problems systematically.

- **BH uniqueness theorem** [Robinson; Mazur; Carter; Israel; Amsel, Horowitz, Marolf, and Roberts]: In 4D, a stationary asymptotically flat black hole is Kerr-Newman.
- The theorem breaks down in higher dimensions, e.g. in 5D, Myers–Perry solutions [Myers and Perry] and black ring solutions [Empanan and Reall] have the same set of conserved charges.
- **Deformation of near-horizon geometry** [Li and Lucietti; Katona and Lucietti; Fontanella and Gutowski]: Finiteness of linear transverse deformations. A black hole with a static and maximally symmetric near-horizon geometry is Schwarzschild–de Sitter.
- L_∞ -**algebra** (review [Jurčo, Raspollini, Sämann, and Wolf]) is a powerful framework in classical and quantum field theory. The **homological perturbation lemma** allows one to solve deformation problems systematically.

- Gaussian null coordinates and near-horizon geometries
- Deformations of near-horizon geometries
- L_∞ -algebras, quasi-isomorphisms and minimal models
- Solving the deformation problem
- Summary and outlook

Gaussian Null Coordinates and Near-horizon Geometries

Gaussian null coordinates and extremal black holes

An **extremal black hole** is a black hole with a $(d - 1)$ -dimensional **Killing horizon** Σ (null hypersurface generated by a Killing vector) such that surface gravity vanishes, i.e.

$$d(K \cdot K)|_{\Sigma} = 0 ,$$

where K is the timelike Killing vector.

Theorem ([Moncrief and Isenberg])

*Near this horizon, there exist **Gaussian Null coordinates**, such that the metric takes the form*

$$g = du \odot [dr + r\alpha_i(r, y)dy^i - \frac{1}{2}r^2\beta(r, y)du] + \frac{1}{2}\gamma_{ij}(r, y)dy^i \odot dy^j ,$$

*where $K = \partial_u$. The space is foliated by $(d - 2)$ -dimensional submanifolds S parametrised by y^i called **spatial cross-section** (assumed to be compact with no boundary).*

Gaussian null coordinates and extremal black holes

An **extremal black hole** is a black hole with a $(d - 1)$ -dimensional **Killing horizon** Σ (null hypersurface generated by a Killing vector) such that surface gravity vanishes, i.e.

$$d(K \cdot K)|_{\Sigma} = 0 ,$$

where K is the timelike Killing vector.


Theorem ([Moncrief and Isenberg])

*Near this horizon, there exist **Gaussian Null coordinates**, such that the metric takes the form*

$$g = du \odot [dr + r\alpha_i(r, y)dy^i - \frac{1}{2}r^2\beta(r, y)du] + \frac{1}{2}\gamma_{ij}(r, y)dy^i \odot dy^j ,$$

*where $K = \partial_u$. The space is foliated by $(d - 2)$ -dimensional submanifolds S parametrised by y^i called **spatial cross-section** (assumed to be compact with no boundary).*

Gaussian null coordinates and extremal black holes



Theorem ([Moncrief and Isenberg])

Near this horizon, there exist **Gaussian Null coordinates**, such that the metric takes the form

$$g = du \odot \left[dr + r\alpha_i(r, y)dy^i - \frac{1}{2}r^2\beta(r, y)du \right] + \frac{1}{2}\gamma_{ij}(r, y)dy^i \odot dy^j ,$$

where $K = \partial_u$. The space is foliated by $(d - 2)$ -dimensional submanifolds S parametrised by y^i called **spatial cross-section** (assumed to be compact with no boundary).

Near-horizon geometries

Define a family of metrics via a family of diffeomorphisms $(u, r, y^i) \rightarrow (u/\epsilon, r\epsilon, y^i)$ as

$$g_\epsilon := du \odot [dr + r\alpha_i(\epsilon r, y)dy^i - \frac{1}{2}r^2\beta(\epsilon r, y)du] + \frac{1}{2}\gamma_{ij}(\epsilon r, y)dy^i \odot dy^j$$

Definition (near-horizon geometries)

The **near-horizon limit** corresponds to $\epsilon \rightarrow 0$, and the resulting geometry is called the **near-horizon geometry**. The metric is

$$\mathring{g} = du \odot [dr + r\mathring{\alpha}_i(y)dy^i - \frac{1}{2}r^2\mathring{\beta}(y)du] + \frac{1}{2}\mathring{\gamma}_{ij}(y)dy^i \odot dy^j ,$$

where $(\mathring{\alpha}(y^i), \mathring{\beta}(y^i), \mathring{\gamma}(y^i)) := (\alpha(0, y^i), \beta(0, y^i), \gamma(0, y^i))$.

Near-horizon geometries

Define a family of metrics via a family of diffeomorphisms $(u, r, y^i) \rightarrow (u/\epsilon, r\epsilon, y^i)$ as

$$g_\epsilon := du \odot [dr + r\alpha_i(\epsilon r, y)dy^i - \frac{1}{2}r^2\beta(\epsilon r, y)du] + \frac{1}{2}\gamma_{ij}(\epsilon r, y)dy^i \odot dy^j$$

Definition (near-horizon geometries)

The **near-horizon limit** corresponds to $\epsilon \rightarrow 0$, and the resulting geometry is called the **near-horizon geometry**. The metric is

$$\mathring{g} = du \odot [dr + r\mathring{\alpha}_i(y)dy^i - \frac{1}{2}r^2\mathring{\beta}(y)du] + \frac{1}{2}\mathring{\gamma}_{ij}(y)dy^i \odot dy^j ,$$

where $(\mathring{\alpha}(y^i), \mathring{\beta}(y^i), \mathring{\gamma}(y^i)) := (\alpha(0, y^i), \beta(0, y^i), \gamma(0, y^i))$.

Examples

The Gaussian null coordinates are analogous to the **Eddington–Finkelstein coordinates**. The Schwarzschild solution:

$$g_S = du \odot \left[dr - \left(\frac{1}{2} - \frac{m}{r} \right) du \right] + r^2 d\Omega^2 .$$

The Kerr black hole is **extremal** when the angular momentum is $J = m^2$. Its near-horizon geometry is called **extremal Kerr horizon**:

$$\begin{aligned} \dot{g}_{eK} = & du \odot \left[dr + r \frac{4(1-x^2)}{(1+x^2)^2} d\varphi - r \frac{2x}{1+x^2} dx + r^2 \frac{3-6x^2-x^4}{2m^2(1+x^2)^3} du \right] \\ & + \frac{m^2(1+x^2)}{2(1-x^2)} dx \odot dx + 2m^2 \frac{1-x^2}{1+x^2} d\varphi \odot d\varphi , \end{aligned}$$

where $x \in (-1, 1)$ and $\varphi \in (0, 2\pi)$.

The Gaussian null coordinates are analogous to the **Eddington–Finkelstein coordinates**. The Schwarzschild solution:

$$g_S = du \odot \left[dr - \left(\frac{1}{2} - \frac{m}{r} \right) du \right] + r^2 d\Omega^2 .$$

The Kerr black hole is **extremal** when the angular momentum is $J = m^2$. Its near-horizon geometry is called **extremal Kerr horizon**:

$$\begin{aligned} \dot{g}_{eK} = & du \odot \left[dr + r \frac{4(1-x^2)}{(1+x^2)^2} d\varphi - r \frac{2x}{1+x^2} dx + r^2 \frac{3-6x^2-x^4}{2m^2(1+x^2)^3} du \right] \\ & + \frac{m^2(1+x^2)}{2(1-x^2)} dx \odot dx + 2m^2 \frac{1-x^2}{1+x^2} d\varphi \odot d\varphi , \end{aligned}$$

where $x \in (-1, 1)$ and $\varphi \in (0, 2\pi)$.

Deformations of Near-horizon Geometries

Deformations of near-horizon geometries

Since **near-horizon geometries** arise as the $\epsilon \rightarrow 0$ limit of the family of diffeomorphisms, they satisfy the **Einstein field equations** and can be used as **background metrics**. We perturb these metrics to

$$g := du \odot [dr + r\alpha_i(r, y)dy^i - \frac{1}{2}r^2\beta(r, y)du] + \frac{1}{2}\gamma_{ij}(r, y)dy^i \odot dy^j ,$$

where

$$\alpha_i(r, y) := \hat{\alpha}_i(y) + h_i(r, y) ,$$

$$\beta(r, y) := \hat{\beta}(y) + h(r, y) ,$$

$$\gamma_{ij}(r, y) := \hat{\gamma}_{ij}(y) + h_{ij}(r, y)$$

with deformations h_i , h , and h_{ij} satisfying

$$h_i|_{r=0} = 0 , \quad h|_{r=0} = 0 , \quad \text{and} \quad h_{ij}|_{r=0} = 0 .$$

Deformations of near-horizon geometries

Since **near-horizon geometries** arise as the $\epsilon \rightarrow 0$ limit of the family of diffeomorphisms, they satisfy the **Einstein field equations** and can be used as **background metrics**. We perturb these metrics to

$$g := du \odot [dr + r\alpha_i(r, y)dy^i - \frac{1}{2}r^2\beta(r, y)du] + \frac{1}{2}\gamma_{ij}(r, y)dy^i \odot dy^j ,$$

where

$$\alpha_i(r, y) := \hat{\alpha}_i(y) + h_i(r, y) ,$$

$$\beta(r, y) := \hat{\beta}(y) + h(r, y) ,$$

$$\gamma_{ij}(r, y) := \hat{\gamma}_{ij}(y) + h_{ij}(r, y)$$

with deformations h_i , h , and h_{ij} satisfying

$$h_i|_{r=0} = 0 , \quad h|_{r=0} = 0 , \quad \text{and} \quad h_{ij}|_{r=0} = 0 .$$

Null orthonormal basis

We define an **unperturbed null orthonormal basis** as

$$\mathring{e}^+ := du, \quad \mathring{e}^- := dr + r\mathring{\alpha}_i dy^i - \frac{1}{2}r^2\mathring{\beta} du, \quad \text{and} \quad \mathring{e}^a := dy^i \mathring{e}_i^a,$$

where

$$\mathring{e}_i^a \mathring{e}_j^b \delta_{ab} = \mathring{\gamma}_{ij}.$$

On this basis, the **near-horizon metric** is

$$\mathring{g} = \mathring{e}^+ \odot \mathring{e}^- + \frac{1}{2}\delta_{ab}\mathring{e}^a \odot \mathring{e}^b,$$

and the **perturbed metric** is

$$g = \mathring{g} + \mathring{e}^+ \odot [rh_a \mathring{e}^a - \frac{1}{2}r^2 h \mathring{e}^+] + \frac{1}{2}h_{ab} \mathring{e}^a \odot \mathring{e}^b,$$

where $h_a := \mathring{E}_a^i h_i$, etc. with \mathring{E}_a^i the inverse of \mathring{e}_i^a .

Contracted Bianchi identities and gauge transformations

Theorem

By the **contracted Bianchi identities**, it is enough to restrict ourselves to

$$0 = G^{++}, \quad 0 = G^{a+}, \quad \text{and} \quad 0 = G^{ab} + \Lambda g^{ab},$$

where G is the **Einstein tensor**. The remaining Einstein field equations do not impose any further constraints.

Theorem ([Fontanella and Gutowski])

Residual gauge transformations are fixed by the gauge-fixing condition

$$\delta^{ab} \partial_r h_{ab}|_{r=0} = \Gamma,$$

where $\Gamma(y^i)$ is the unique (up to a multiplicative constant) solution of

$$\overset{\circ}{\nabla}_a \overset{\circ}{\nabla}^a \Gamma + \overset{\circ}{\alpha}^a \overset{\circ}{\nabla}_a \Gamma + \overset{\circ}{\nabla}_a \overset{\circ}{\alpha}^a \Gamma = 0,$$

and $\overset{\circ}{\nabla}$ is the **Levi-Civita connection** on the spatial cross-section \mathring{S} with respect to the metric $\overset{\circ}{\gamma}$.

Contracted Bianchi identities and gauge transformations

Theorem

By the **contracted Bianchi identities**, it is enough to restrict ourselves to

$$0 = G^{++}, \quad 0 = G^{a+}, \quad \text{and} \quad 0 = G^{ab} + \Lambda g^{ab},$$

where G is the **Einstein tensor**. The remaining Einstein field equations do not impose any further constraints.

Theorem ([Fontanella and Gutowski])

Residual gauge transformations are fixed by the **gauge-fixing condition**

$$\delta^{ab} \partial_r h_{ab}|_{r=0} = \Gamma,$$


where $\Gamma(y^i)$ is the unique (up to a multiplicative constant) solution of

$$\overset{\circ}{\nabla}_a \overset{\circ}{\nabla}^a \Gamma + \overset{\circ}{\alpha}^a \overset{\circ}{\nabla}_a \Gamma + \overset{\circ}{\nabla}_a \overset{\circ}{\alpha}^a \Gamma = 0,$$

and $\overset{\circ}{\nabla}$ is the **Levi-Civita connection** on the **spatial cross-section** \mathring{S} with respect to the metric $\overset{\circ}{\gamma}$.

L_∞ -algebras and Minimal Models

Why L_∞ -algebras?



Definition (L_∞ -algebra)

An L_∞ -algebra (V, μ_i) is a \mathbb{Z} -graded vector space $V = \bigoplus_{k \in \mathbb{Z}} V_k$ equipped with **multi-linear maps** $\mu_i : \bigwedge^i V \rightarrow V$ satisfying the **homotopy Jacobi identities**. The first few identities are

$$\begin{aligned}\mu_1(\mu_1(v_1)) &= 0 \\ \mu_1(\mu_2(v_1, v_2)) &= \pm \mu_2(\mu_1(v_1), v_2) \pm \mu_2(\mu_1(v_2), v_1) \\ &\quad \mu_2(\mu_2(v_1, v_2), v_3) \pm \text{cyclic} \\ &= \mu_1(\mu_3(v_1, v_2, v_3)) + \mu_3(\mu_1(v_1), v_2, v_3) \pm \text{cyclic} \\ &\quad \vdots\end{aligned}$$

for homogeneous $v_i \in V$.

Notice that the first identity implies $\mu_1^2 = 0$. This allows us to define a **cochain complex** (V, μ_1) and its **cohomology** $H^\bullet(V)$.

Maurer–Cartan equations

Definition (curvature)

The **curvature** f is a degree 2 element, defined by

$$f := \sum_{i \geq 1} \frac{1}{i!} \mu_i(a, \dots, a),$$

where a is a degree one element in an L_∞ -algebra (V, μ_i)

Definition (Maurer–Cartan equation)

The **Maurer–Cartan equation** is defined by the vanishing of the curvature:

$$\sum_{i \geq 1} \frac{1}{i!} \mu_i(a, \dots, a) = 0.$$

Maurer–Cartan equations

Definition (curvature)

The **curvature** f is a degree 2 element, defined by

$$f := \sum_{i \geq 1} \frac{1}{i!} \mu_i(a, \dots, a),$$

where a is a degree one element in an L_∞ -algebra (V, μ_i)

Definition (Maurer–Cartan equation)

The **Maurer–Cartan equation** is defined by the vanishing of the curvature:

$$\sum_{i \geq 1} \frac{1}{i!} \mu_i(a, \dots, a) = 0.$$

Deformation of near-horizon geometry

For our purpose, our L_∞ -algebra is concentrated in **degrees 1 and 2**, that is, $V = V_1 \oplus V_2$

$$\Theta := (h_a, h, h_{ab})$$
$$V_1 := \Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)|_{\text{gf}}$$

$$(G^{a+}, G^{++}, G^{ab} + \Lambda g^{ab})$$
$$V_2 := \Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)$$

The **Maurer–Cartan equation** is

$$\mu_1(\Theta) + \frac{1}{2}\mu_2(\Theta, \Theta) + \dots = 0,$$

where $\mu_1(\Theta)$ is the **linear part** of the Einstein field equations, $\mu_2(\Theta, \Theta)$ is the **bilinear part**, and so on. μ_i vanishes if any argument is in V_2 .

Deformation of near-horizon geometry

For our purpose, our L_∞ -algebra is concentrated in **degrees 1 and 2**, that is, $V = V_1 \oplus V_2$

$$\begin{aligned} \Theta &:= (h_a, h, h_{ab}) \\ V_1 &:= \Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)|_{\text{gf}} \\ &\quad (G^{a+}, G^{++}, G^{ab} + \Lambda g^{ab}) \\ V_2 &:= \Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S) \end{aligned}$$

The **Maurer–Cartan equation** is

$$\mu_1(\Theta) + \frac{1}{2}\mu_2(\Theta, \Theta) + \dots = 0,$$

where $\mu_1(\Theta)$ is the **linear part** of the Einstein field equations, $\mu_2(\Theta, \Theta)$ is the **bilinear part**, and so on. μ_i vanishes if any argument is in V_2 .

Definition (L_∞ -algebra morphism)

L_∞ -algebra morphism from (V, μ_i) to (V', μ'_i) is a set of maps $\phi_i : \bigwedge^i V \rightarrow V'$ satisfying the **compatibility relations**. The first few are

$$\begin{aligned}\phi_1(\mu_1(v_1)) &= \mu'_1(\phi_1(v_1)) , \\ \phi_1(\mu_2(v_1, v_2)) &- \mu'_2(\phi_1(v_1), \phi_1(v_2)) \\ &= \pm \phi_2(\mu_1(v_1), v_2) \pm \phi_2(\mu_1(v_2), v_1) \pm \mu'_1(\phi_2(v_1, v_2)) \\ &\quad \vdots\end{aligned}$$

Definition (quasi-isomorphism)

A **quasi-isomorphism** is an L_∞ -algebra morphism such that ϕ_1 induces an **isomorphism between cohomologies** $H^\bullet(V)$ and $H^\bullet(V')$ of the underlying cochain complexes (V, μ_1) and (V', μ'_1) .

Theorem

If two L_∞ -algebras are quasi-isomorphic, the moduli spaces of solutions (modulo gauge) of their Maurer–Cartan equations are isomorphic. The isomorphism is given by

$$a' = \sum_{i \geq 1} \frac{1}{i!} \phi_i(a, \dots, a)$$

for $a \in V_1$ and $a' \in V'_1$

Definition (quasi-isomorphism)

A **quasi-isomorphism** is an L_∞ -algebra morphism such that ϕ_1 induces an **isomorphism between cohomologies** $H^\bullet(V)$ and $H^\bullet(V')$ of the underlying cochain complexes (V, μ_1) and (V', μ'_1) .

Theorem

If two L_∞ -algebras are **quasi-isomorphic**, the **moduli spaces of solutions (modulo gauge) of their Maurer–Cartan equations are isomorphic**. The isomorphism is given by

$$a' = \sum_{i \geq 1} \frac{1}{i!} \phi_i(a, \dots, a)$$

for $a \in V_1$ and $a' \in V'_1$

Theorem

Given **cochain complexes** (V, μ_1) and $(H^\bullet(V), 0)$ along with **contracting homotopy** h and **cochain maps** p and e

$$h \circlearrowleft (V, \mu_1) \begin{matrix} \xrightarrow{p} \\ \xleftarrow{e} \end{matrix} (H^\bullet(V), 0) ,$$

satisfying

$$\begin{aligned} id &= \mu_1 \circ h + h \circ \mu_1 + e \circ p, \\ p \circ e &= id, \quad p \circ \mu_1 = 0, \quad \mu_1 \circ e = 0, \\ p \circ h &= h \circ e = h \circ h = 0. \end{aligned}$$

There exists an L_∞ -algebra $(H^\bullet(V), \mu_i^\circ)$ with $\mu_1^\circ = 0$, quasi-isomorphic to (V, μ_i) . The **quasi-isomorphism** is $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$ and $E_1 = e$.

Theorem (contd.)

As an example,

$$\begin{aligned} E_2(v_1^\circ, v_2^\circ) &:= -h\{\mu_2[E_1(v_1^\circ), E_1(v_2^\circ)]\} , \\ E_3(v_1^\circ, v_2^\circ, v_3^\circ) &:= \pm h\{\mu_3[E_1(v_1^\circ), E_1(v_2^\circ), E_1(v_3^\circ)] \\ &\quad \pm \mu_2[E_2(v_1^\circ, v_2^\circ), E_1(v_3^\circ)] \pm \text{cyclic}\} , \\ &\vdots \\ \mu_2^\circ(v_1^\circ, v_2^\circ) &:= p\{\mu_2[E_1(v_1^\circ), E_1(v_2^\circ)]\} , \\ \mu_3^\circ(v_1^\circ, v_2^\circ, v_3^\circ) &:= \pm p\{\mu_3[E_1(v_1^\circ), E_1(v_2^\circ), E_1(v_3^\circ)] \\ &\quad \pm \mu_2[E_2(v_1^\circ, v_2^\circ), E_1(v_3^\circ)] \pm \text{cyclic}\} , \\ &\vdots \end{aligned}$$

for all homogeneous $v_1^\circ, \dots, v_i^\circ \in H^\bullet(V)$.

Minimal model contd.

Maps h , p and e and their conditions are the **abstract Hodge decomposition**

$$\begin{array}{ccc}
 \vdots & & \vdots \\
 V^{k-1} = H^{k-1} \oplus B^{k-1} \oplus I^{k-1} & & H^{k-1}(V) \\
 & \swarrow \mu_1 & \\
 & & \nearrow h_k \\
 V^k = H^k \oplus B^k \oplus I^k & & H^k(V) \\
 & \searrow e_k & \\
 \vdots & & \vdots
 \end{array}$$

μ_1 (arrow from H^k to H^{k-1})
 h_k (arrow from H^{k-1} to H^k)
 e_k (arrow from $H^k(V)$ to H^k)
 p_k (arrow from $H^k(V)$ to H^k)

where H^k , B^k and I^k are the **harmonic**, **exact** and **coexact** subspaces of V_k . p_k and h_k are the inverses of μ_1 and e_k on the restricted spaces, which are trivially extended to all of V_k .

Deformation of near-horizon geometry

In our case, $V = V_1 \oplus V_2$ we have

$$\begin{array}{ccccc}
 & V_1 & \xrightarrow{\mu_1} & V_2 & \\
 \text{id}|_{V_1 - \epsilon_1 \circ \pi_1} \nearrow & & & & \searrow \text{id}|_{V_2 - \epsilon_2 \circ \pi_2} \\
 & \swarrow \iota_1 & & \swarrow \iota_2 & \\
 \ker(\mu_1) & & & & \text{im}(\mu_1) \\
 = H^1(V) & & \xrightarrow{\hat{\mu}_1} & & \\
 & \searrow \epsilon_1 & & \searrow \epsilon_2 & \\
 & V_1 / \ker(\mu_1) & & V_2 / \text{im}(\mu_1) & \\
 & & & = H^2(V) &
 \end{array}$$

with $\pi_{1,2}$ the **canonical quotient projections** and $\epsilon_{1,2}$ choices of **right-inverses**, $\iota_{1,2}$ the **inclusions**, and $\hat{\mu}_1$ the **canonical isomorphism** given by the first isomorphism theorem.

We define

$$h \circlearrowleft (V, \mu_1) \xrightleftharpoons[e]{p} (H^\bullet(V), 0),$$

where

$$\begin{aligned}
 p|_{V_1} &:= \text{id}|_{V_1 - \epsilon_1 \circ \pi_1} = \text{id}|_{V_1} - h \circ \mu_1, & p|_{V_2} &:= \pi_2, \\
 e|_{H^1(V)} &:= \iota_1, & e|_{H^2(V)} &:= \epsilon_2, & h &:= \epsilon_1 \circ \hat{\mu}_1^{-1} \circ (\text{id}|_{V_2 - \epsilon_2 \circ \pi_2}).
 \end{aligned}$$

Deformation of near-horizon geometry

In our case, $V = V_1 \oplus V_2$ we have

$$\begin{array}{ccccc}
 & V_1 & \xrightarrow{\mu_1} & V_2 & \\
 \text{id}|_{V_1} - \epsilon_1 \circ \pi_1 \nearrow & & & & \searrow \text{id}|_{V_2} - \epsilon_2 \circ \pi_2 \\
 & \swarrow \iota_1 & & \swarrow \iota_2 & \\
 \ker(\mu_1) & & & & \text{im}(\mu_1) \\
 = H^1(V) & & \xrightarrow{\hat{\mu}_1} & & \\
 & \searrow \epsilon_1 & & \searrow \epsilon_2 & \\
 & V_1 / \ker(\mu_1) & & V_2 / \text{im}(\mu_1) & \\
 & & & = H^2(V) &
 \end{array}$$

with $\pi_{1,2}$ the **canonical quotient projections** and $\epsilon_{1,2}$ choices of **right-inverses**, $\iota_{1,2}$ the **inclusions**, and $\hat{\mu}_1$ the **canonical isomorphism** given by the first isomorphism theorem.

We define

$$\text{h} \circlearrowleft (V, \mu_1) \xrightleftharpoons[e]{\text{p}} (H^\bullet(V), 0),$$

where

$$\begin{aligned}
 \text{p}|_{V_1} &:= \text{id}|_{V_1} - \epsilon_1 \circ \pi_1 = \text{id}|_{V_1} - \text{h} \circ \mu_1, & \text{p}|_{V_2} &:= \pi_2, \\
 \text{e}|_{H^1(V)} &:= \iota_1, & \text{e}|_{H^2(V)} &:= \epsilon_2, & \text{h} &:= \epsilon_1 \circ \hat{\mu}_1^{-1} \circ (\text{id}|_{V_2} - \epsilon_2 \circ \pi_2).
 \end{aligned}$$

Solving the deformation problem

Outline of calculation steps

- Construct an L_∞ -**algebra** (V, μ_i) by reading off the μ_i from the **Einstein field equations**.
- Construct the **cochain complex** (V, μ_1) and identify its **cohomology**.
- Determine the **contracting homotopy** h in

$$h \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} (V, \mu_1) \begin{array}{c} \xleftarrow{p} \\ \xrightarrow{e} \end{array} \gg (H^\bullet(V), 0).$$

- Construct the **minimal model** $(H^\bullet(V), \mu_i^\circ)$ and the **quasi-isomorphism** $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$
- Solve the **Maurer–Cartan equation** of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- Use the **quasi-isomorphism** E to transform these solutions to solutions of the Maurer–Cartan equation of (V, μ_i)

Outline of calculation steps

- Construct an L_∞ -**algebra** (V, μ_i) by reading off the μ_i from the **Einstein field equations**.
- Construct the **cochain complex** (V, μ_1) and identify its **cohomology**.
- Determine the **contracting homotopy** h in

$$h \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} (V, \mu_1) \begin{array}{c} \xleftarrow{p} \\ \xrightarrow{e} \end{array} \gg (H^\bullet(V), 0).$$

- Construct the **minimal model** $(H^\bullet(V), \mu_i^\circ)$ and the **quasi-isomorphism** $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$
- Solve the **Maurer–Cartan equation** of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- Use the **quasi-isomorphism** E to transform these solutions to solutions of the Maurer–Cartan equation of (V, μ_i)

Outline of calculation steps

- Construct an L_∞ -**algebra** (V, μ_i) by reading off the μ_i from the **Einstein field equations**.
- Construct the **cochain complex** (V, μ_1) and identify its **cohomology**.
- Determine the **contracting homotopy** h in

$$h \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} (V, \mu_1) \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{e} \end{array} (H^\bullet(V), 0).$$

- Construct the **minimal model** $(H^\bullet(V), \mu_i^\circ)$ and the **quasi-isomorphism** $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$
- Solve the **Maurer–Cartan equation** of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- Use the **quasi-isomorphism** E to transform these solutions to solutions of the Maurer–Cartan equation of (V, μ_i)

Outline of calculation steps

- Construct an L_∞ -**algebra** (V, μ_i) by reading off the μ_i from the **Einstein field equations**.
- Construct the **cochain complex** (V, μ_1) and identify its **cohomology**.
- Determine the **contracting homotopy** h in

$$h \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} (V, \mu_1) \begin{array}{c} \xleftarrow{p} \\ \xrightarrow{e} \end{array} \gg (H^\bullet(V), 0).$$

- Construct the **minimal model** $(H^\bullet(V), \mu_i^\circ)$ and the **quasi-isomorphism** $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$
- Solve the **Maurer–Cartan equation** of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- Use the **quasi-isomorphism** E to transform these solutions to solutions of the Maurer–Cartan equation of (V, μ_i)

Outline of calculation steps

- Construct an L_∞ -**algebra** (V, μ_i) by reading off the μ_i from the **Einstein field equations**.
- Construct the **cochain complex** (V, μ_1) and identify its **cohomology**.
- Determine the **contracting homotopy** h in

$$h \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} (V, \mu_1) \begin{array}{c} \xleftarrow{p} \\ \xrightarrow{e} \end{array} \gg (H^\bullet(V), 0).$$

- Construct the **minimal model** $(H^\bullet(V), \mu_i^\circ)$ and the **quasi-isomorphism** $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$
- Solve the **Maurer–Cartan equation** of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- Use the **quasi-isomorphism** E to transform these solutions to solutions of the Maurer–Cartan equation of (V, μ_i)

Outline of calculation steps

- Construct an L_∞ -**algebra** (V, μ_i) by reading off the μ_i from the **Einstein field equations**.
- Construct the **cochain complex** (V, μ_1) and identify its **cohomology**.
- Determine the **contracting homotopy** h in

$$h \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} (V, \mu_1) \begin{array}{c} \xleftarrow{p} \\ \xrightarrow{e} \end{array} \gg (H^\bullet(V), 0).$$

- Construct the **minimal model** $(H^\bullet(V), \mu_i^\circ)$ and the **quasi-isomorphism** $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$
- Solve the **Maurer–Cartan equation** of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- Use the **quasi-isomorphism** E to transform these solutions to solutions of the Maurer–Cartan equation of (V, μ_i)

Outline of calculation steps

- Construct an L_∞ -**algebra** (V, μ_i) by reading off the μ_i from the **Einstein field equations**.
- Construct the **cochain complex** (V, μ_1) and **identify its cohomology**.
- Determine the **contracting homotopy** h in

$$h \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} (V, \mu_1) \begin{array}{c} \xleftarrow{p} \\ \xrightarrow{e} \end{array} \gg (H^\bullet(V), 0).$$

- Construct the **minimal model** $(H^\bullet(V), \mu_i^\circ)$ and the **quasi-isomorphism** $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$
- Solve the **Maurer–Cartan equation** of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- Use the **quasi-isomorphism** E to transform these solutions to solutions of the Maurer–Cartan equation of (V, μ_i)

Cochain complex and its cohomology

The **cochain complex** is

$$\underbrace{\Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)}_{=: V_1} \Big|_{\text{gf}} \xrightarrow{\mu_1} \underbrace{\Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)}_{=: V_2} .$$

We find that

$$H^1(V) = \ker(\mu_1) \cong \ker(\bar{d}_{ab}{}^{cd})$$

with

$$\bar{d}_{ab}{}^{cd} := r^2 \bar{a}_{ab}{}^{cd}(y) \partial_r^2 + r \bar{b}_{ab}{}^{cd}(y) \partial_r + \bar{c}_{ab}{}^{cd}(y) ,$$

where $\bar{a}_{ab}{}^{cd}$, $\bar{b}_{ab}{}^{cd}$ and $\bar{c}_{ab}{}^{cd}$ are **operators on the spatial cross-section** S mapping the traceless part of h_{cd} (denoted \bar{h}_{cd}) to traceless symmetric $(0, 2)$ -tensors.

Expanding $\bar{h}_{cd}(r, y) = \sum_{n>0} \frac{r^n}{n!} \bar{h}_{cd}^{(n)}(y)$, the space of solutions of

$$[n(n-1)\bar{a}_{ab}{}^{cd}(y) + n\bar{b}_{ab}{}^{cd}(y) + \bar{c}_{ab}{}^{cd}(y)] \bar{h}_{cd}^{(n)}(y) = 0$$

is **isomorphic** to $H^1(V)$.

Cochain complex and its cohomology

The **cochain complex** is

$$\underbrace{\Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)}_{=: V_1} \Big|_{\text{gf}} \xrightarrow{\mu_1} \underbrace{\Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)}_{=: V_2} .$$

We find that

$$H^1(V) = \ker(\mu_1) \cong \ker(\bar{d}_{ab}{}^{cd})$$

with

$$\bar{d}_{ab}{}^{cd} := r^2 \bar{a}_{ab}{}^{cd}(y) \partial_r^2 + r \bar{b}_{ab}{}^{cd}(y) \partial_r + \bar{c}_{ab}{}^{cd}(y) ,$$

where $\bar{a}_{ab}{}^{cd}$, $\bar{b}_{ab}{}^{cd}$ and $\bar{c}_{ab}{}^{cd}$ are **operators on the spatial cross-section** S mapping the traceless part of h_{cd} (denoted \bar{h}_{cd}) to traceless symmetric $(0, 2)$ -tensors.

Expanding $\bar{h}_{cd}(r, y) = \sum_{n>0} \frac{r^n}{n!} \bar{h}_{cd}^{(n)}(y)$, the space of solutions of

$$[n(n-1)\bar{a}_{ab}{}^{cd}(y) + n\bar{b}_{ab}{}^{cd}(y) + \bar{c}_{ab}{}^{cd}(y)] \bar{h}_{cd}^{(n)}(y) = 0$$

is **isomorphic** to $H^1(V)$.

Cochain complex and its cohomology

The **cochain complex** is

$$\underbrace{\Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)}_{=: V_1} \Big|_{\text{gf}} \xrightarrow{\mu_1} \underbrace{\Omega_r^1(S) \oplus \mathcal{C}_r^\infty(S) \oplus \mathcal{S}_r^2(S)}_{=: V_2} .$$

We find that

$$H^1(V) = \ker(\mu_1) \cong \ker(\bar{d}_{ab}{}^{cd})$$

with

$$\bar{d}_{ab}{}^{cd} := r^2 \bar{a}_{ab}{}^{cd}(y) \partial_r^2 + r \bar{b}_{ab}{}^{cd}(y) \partial_r + \bar{c}_{ab}{}^{cd}(y) ,$$

where $\bar{a}_{ab}{}^{cd}$, $\bar{b}_{ab}{}^{cd}$ and $\bar{c}_{ab}{}^{cd}$ are **operators on the spatial cross-section** S mapping the traceless part of h_{cd} (denoted \bar{h}_{cd}) to traceless symmetric $(0, 2)$ -tensors.

Expanding $\bar{h}_{cd}(r, y) = \sum_{n>0} \frac{r^n}{n!} \bar{h}_{cd}^{(n)}(y)$, the space of solutions of

$$[n(n-1)\bar{a}_{ab}{}^{cd}(y) + n\bar{b}_{ab}{}^{cd}(y) + \bar{c}_{ab}{}^{cd}(y)] \bar{h}_{cd}^{(n)}(y) = 0$$

is **isomorphic** to $H^1(V)$.

Cochain complex and its cohomology contd.

$$[n(n-1)\bar{a}_{ab}{}^{cd}(y) + n\bar{b}_{ab}{}^{cd}(y)\partial_r + \bar{c}_{ab}{}^{cd}(y)]h_{cd}^{(n)}(y) = 0$$

$$\bar{a}_{ab}{}^{cd} := (\delta_{(a}{}^c\delta_{b)}{}^d - \frac{1}{d-2}\delta_{ab}\delta^{cd})\left(\frac{1}{4}\dot{\alpha}^e\dot{\alpha}_e + \frac{1}{4}\dot{\nabla}^e\dot{\alpha}_e - \frac{1}{d-2}\Lambda\right),$$

$$\begin{aligned} \bar{b}_{ab}{}^{cd} := & \delta_{(b}{}^{(c}[\dot{\nabla}^d)\dot{\alpha}_a] - \dot{\nabla}_a\dot{\alpha}^d) + \delta_a{}^d)\left(\frac{1}{2}\dot{\alpha}_e\dot{\alpha}^e - \dot{\alpha}^e\dot{\nabla}_e - \frac{2}{d-2}\Lambda\right) \\ & - \frac{1}{d-2}\delta_{ab}\delta^{cd}\left(\frac{1}{2}\dot{\alpha}_e\dot{\alpha}^e - \dot{\alpha}^e\dot{\nabla}_e - \frac{2}{d-2}\Lambda\right), \end{aligned}$$

$$\begin{aligned} \bar{c}_{ab}{}^{cd} := & \delta_{(b}{}^{(c}\{\dot{\alpha}^d)\dot{\nabla}_a] - \dot{\alpha}_a)\dot{\nabla}^d) - \dot{\nabla}^d)\dot{\nabla}_a] + \dot{\nabla}_a)\dot{\nabla}^d) \\ & + \delta_a{}^d)\left[\frac{1}{2}(\dot{\nabla}_e - \dot{\alpha}_e)\dot{\nabla}^e + \frac{2}{d-2}\Lambda\right]\} \\ & - \frac{1}{d-2}\delta_{ab}\delta^{cd}\left[\frac{1}{2}(\dot{\nabla}_e - \dot{\alpha}_e)\dot{\nabla}^e + \frac{2}{d-2}\Lambda\right]. \end{aligned}$$

For each n , the equation is **elliptic** on S (compact without boundary). Thus, at each order in r , $H^1(V)$ is **finite dimensional**, which implies that the **moduli space of all-order solutions** (smaller than $H^1(V)$) is **finite dimensional** at each order in r .

Example (extremal Kerr horizon)

$$[n(n-1)\bar{a}_{ab}{}^{cd}(y) + n\bar{b}_{ab}{}^{cd}(y)\partial_r + \bar{c}_{ab}{}^{cd}(y)]h_{cd}^{(n)}(y) = 0$$

are difficult to solve for arbitrary near-horizon geometries. However, for **extremal Kerr near-horizon geometry**, assuming **axisymmetry**, the solutions are

$$\begin{aligned}\bar{h}_{11}^{(1)}(x) &= \frac{A(1-x^2)(5-16x^2-5x^4)}{m(1+x^2)^2}, \\ \bar{h}_{12}^{(1)}(x) &= \frac{A x(1-x^2)(9+x^2)}{m(1+x^2)^2}, \\ \bar{h}_{11}^{(n)}(x) &= \frac{(K_1^{(n)}(1-x^2) - 2K_3^{(n)}x)P_n^2(x)}{m^n(1+x^2)^{n+1}}, \\ \bar{h}_{12}^{(n)} &= \frac{(2K_1^{(n)}x + K_3^{(n)}(1-x^2))P_n^2(x)}{m^n(1+x^2)^{n+1}}\end{aligned}$$

for $n > 1$, where A , $K_1^{(n)}$ and $K_3^{(n)}$ are arbitrary **constants** and $P_n^2(x)$ denotes the associated Legendre polynomial.

Contracting homotopy h

The **contracting homotopy** h is uniquely determined by the **Green function** of $\bar{d}_{ab}{}^{cd}$, i.e.

$$\begin{aligned} & [n(n-1)\bar{a}_{ab}{}^{cd}(y) + n\bar{b}_{ab}{}^{cd}(y) + \bar{c}_{ab}{}^{cd}(y)]\bar{g}_{ab}{}^{(n)cd}(y; y') \\ &= \left(\delta_{(a}{}^c\delta_{b)}{}^d - \frac{1}{d-2}\delta_{ab}\delta^{cd}\right)\frac{\delta^{(d-2)}(y-y')}{\sqrt{\det(\hat{\gamma}(y))}}. \end{aligned}$$

The relation between h and $g_{ab}{}^{(n)cd}$ is lengthy and will be omitted. For **extremal Kerr near-horizon geometry**, the Green function is

$$\begin{aligned} \bar{g}_{11}^{(1)11}(x, \varphi; x', \varphi') &= -\frac{(1+x'^2)[1-x^2+4xx'(1-x^2)]}{24(1-x^2)(1-x'^2)(1+x^2)^2}\delta(\varphi-\varphi') \\ &\quad \times [(2-x)(1+x)^2(2+x')(1-x')^2\theta(x'-x) \\ &\quad + (1-x)^2(2+x)(2-x')(1+x')^2\theta(x-x')], \\ \bar{g}_{11}^{(n)11}(x, \varphi; x', \varphi') &= \frac{(1+x^2)^{-n-1}(1+x'^2)^n(1-x+x')(1-x'+x(1+x'))}{2(n-1)n(n+1)(n+2)} \\ &\quad \times [\theta(x-x')P_n^2(x')Q_n^2(x) \\ &\quad + \theta(x'-x)P_n^2(x)Q_n^2(x')]\delta(\varphi-\varphi'), \\ \bar{g}_{22}^{(1)12}(x, \varphi; x', \varphi') &= -\frac{(1+x^2)(x-x')(1+xx')}{12(1-x^2)(1-x'^2)(1+x^2)^2}\delta(\varphi-\varphi') \\ &\quad \times [(2-x)(1+x)^2(2+x')(1-x')^2\theta(x'-x) \\ &\quad + (1-x)^2(2+x)(2-x')(1+x')^2\theta(x-x')], \\ \bar{g}_{22}^{(n)12}(x, \varphi; x', \varphi') &= \frac{(1+x^2)^{-n-1}(1+x'^2)^n(x-x')(1+xx')}{(n-1)n(n+1)(n+2)} \\ &\quad \times [\theta(x-x')P_n^2(x')Q_n^2(x) \\ &\quad + \theta(x'-x)P_n^2(x)Q_n^2(x')]\delta(\varphi-\varphi') \end{aligned}$$

for $n > 1$ and $\bar{g}_{11}{}^{11} = \bar{g}_{12}{}^{12}$ and $\bar{g}_{11}{}^{12} = -\bar{g}_{12}{}^{11}$

Contracting homotopy h

The **contracting homotopy** h is uniquely determined by the **Green function** of $\bar{d}_{ab}{}^{cd}$, i.e.

$$\begin{aligned} & [n(n-1)\bar{a}_{ab}{}^{cd}(y) + n\bar{b}_{ab}{}^{cd}(y) + \bar{c}_{ab}{}^{cd}(y)]\bar{g}_{ab}{}^{(n)cd}(y; y') \\ &= \left(\delta_{(a}{}^c\delta_{b)}{}^d - \frac{1}{d-2}\delta_{ab}\delta^{cd}\right)\frac{\delta^{(d-2)}(y-y')}{\sqrt{\det(\hat{\gamma}(y))}}. \end{aligned}$$

The relation between h and $\bar{g}_{ab}{}^{(n)cd}$ is lengthy and will be omitted. For **extremal Kerr near-horizon geometry**, the Green function is

$$\begin{aligned} \bar{g}_{11}^{(1)11}(x, \varphi; x', \varphi') &= -\frac{(1+x'^2)[1-x^2+4xx'-x^2(1-x^2)]}{24(1-x^2)(1-x'^2)(1+x^2)^2}\delta(\varphi-\varphi') \\ &\quad \times [(2-x)(1+x)^2(2+x')(1-x')^2\theta(x'-x) \\ &\quad + (1-x)^2(2+x)(2-x')(1+x')^2\theta(x-x')], \\ \bar{g}_{11}^{(n)11}(x, \varphi; x', \varphi') &= \frac{(1+x^2)^{-n-1}(1+x'^2)^n(1-x+x'(1+x))(1-x'+x(1+x'))}{2(n-1)n(n+1)(n+2)} \\ &\quad \times [\theta(x-x')P_n^2(x')Q_n^2(x) \\ &\quad + \theta(x'-x)P_n^2(x)Q_n^2(x')]\delta(\varphi-\varphi'), \\ \bar{g}_{11}^{(1)12}(x, \varphi; x', \varphi') &= -\frac{(1+x^2)(x-x')(1+xx')}{12(1-x^2)(1-x'^2)(1+x^2)^2}\delta(\varphi-\varphi') \\ &\quad \times [(2-x)(1+x)^2(2+x')(1-x')^2\theta(x'-x) \\ &\quad + (1-x)^2(2+x)(2-x')(1+x')^2\theta(x-x')], \\ \bar{g}_{11}^{(n)12}(x, \varphi; x', \varphi') &= \frac{(1+x^2)^{-n-1}(1+x'^2)^n(x-x')(1+xx')}{(n-1)n(n+1)(n+2)} \\ &\quad \times [\theta(x-x')P_n^2(x)Q_n^2(x) \\ &\quad + \theta(x'-x)P_n^2(x)Q_n^2(x')]\delta(\varphi-\varphi') \end{aligned}$$

for $n > 1$ and $\bar{g}_{11}{}^{11} = \bar{g}_{12}{}^{12}$ and $\bar{g}_{11}{}^{12} = -\bar{g}_{12}{}^{11}$

The minimal model

We construct solutions only up to **next-to-lowest order**, so we consider μ_i and E_i for $i \leq 2$.

- The **minimal model** is $(H^\bullet(V) = H^1(V) \oplus H^2(V), \mu_i^\circ)$, where

$$\mu_1^\circ(\Theta_1^\circ) := 0, \quad \mu_2^\circ(\Theta_1^\circ, \Theta_2^\circ) := p(\mu_2(E_1(\Theta_1^\circ), E_1(\Theta_2^\circ))), \dots$$

for all homogeneous $\Theta_1^\circ, \Theta_2^\circ \in H^\bullet(V)$.

- The **quasi-isomorphism** $E : (H^\bullet(V), \mu_i^\circ) \rightarrow (V, \mu_i)$ is

$$E_1 := \text{id}, \quad E_2(\Theta_1^\circ, \Theta_2^\circ) := -h(\mu_2(E_1(\Theta_1^\circ), E_1(\Theta_2^\circ))), \dots$$

- The **Maurer–Cartan equation** of $(H^\bullet(V), \mu_i^\circ)$ is

$$\frac{1}{2}\mu_2^\circ(\Theta^\circ, \Theta^\circ) + \dots = 0$$

for $\Theta^\circ \in H^1(V)$.

Example (extremal Kerr horizon)

The space $H^1(V)$ is parametrised by A , $K_1^{(n)}$ and $K_3^{(n)}$. Hence, the **Maurer–Cartan equation** is algebraic,

$$\text{MC}(A, K_1^{(n)}, K_3^{(n)}) = 0 .$$

At **next-to-lowest order**, this is **trivially satisfied**, so the space of solutions of the **minimal model** Maurer–Cartan equation is $H^1(V)$. We map these solutions to those of the **original** Maurer–Cartan equation of (V, μ_1) (the **Einstein field equations**) via

$$\Theta = E_1(\Theta^\circ) + \frac{1}{2}E_2(\Theta^\circ, \Theta^\circ) + \dots .$$

Hence, the moduli space of **next-to-lowest order deformations** of extremal Kerr near-horizon geometry is parametrised by A , $K_1^{(n)}$ and $K_3^{(n)}$. At **higher orders**, A , $K_1^{(n)}$ and $K_3^{(n)}$ may become constrained.

Example (extremal Kerr horizon)

The space $H^1(V)$ is parametrised by A , $K_1^{(n)}$ and $K_3^{(n)}$. Hence, the **Maurer–Cartan equation** is algebraic,

$$\text{MC}(A, K_1^{(n)}, K_3^{(n)}) = 0 .$$

At **next-to-lowest order**, this is **trivially satisfied**, so the space of solutions of the **minimal model** Maurer–Cartan equation is $H^1(V)$. We map these solutions to those of the **original** Maurer–Cartan equation of (V, μ_1) (the **Einstein field equations**) via

$$\Theta = E_1(\Theta^\circ) + \frac{1}{2}E_2(\Theta^\circ, \Theta^\circ) + \dots .$$

Hence, the moduli space of **next-to-lowest order deformations** of extremal Kerr near-horizon geometry is parametrised by A , $K_1^{(n)}$ and $K_3^{(n)}$. At **higher orders**, A , $K_1^{(n)}$ and $K_3^{(n)}$ may become constrained.

Example (extremal Kerr horizon) contd.

As an example the **next-to-lowest order solutions** for h_{11} are

$$\begin{aligned} h_{11} = & \frac{Ar(5 - 3x^2 + 13x^4 + 5x^6)}{5m(1+x^2)^2} - \frac{3r^2(1-x^2)}{2m^2(1+x^2)^3} \left[-K_1^{(2)}(1-x^2) + 2K_3^{(2)}x \right] \\ & + \frac{A^2r^2}{300m^2(1+x^2)^4} (-675 + 92x + 2752x^2 - 1851x^4 - 92x^5 - 268x^6 + 747x^8 \\ & + 420x^{10} + 75x^{12}) + \mathcal{O}(r^3) + \dots \end{aligned}$$

To locate the **extremal Kerr black hole** in the **moduli space of deformations**, we can either:

- Compare it to the **next-to-lowest order solutions** to fix A , $K_1^{(2)}$ and $K_3^{(2)}$.
- Projects it onto $H^1(V)$ using p_1 , then compare it to the **lowest-order solution** to fix all A , $K_1^{(n)}$ and $K_3^{(n)}$.

We find

$$A = \frac{15}{7}, \quad K_1^{(2)} = \frac{277}{49}, \quad K_3^{(2)} = \frac{23}{49}, \quad K_1^{(3)} = \frac{2684}{1029}, \quad K_3^{(3)} = -\frac{254}{343}.$$

Example (extremal Kerr horizon) contd.

As an example the **next-to-lowest order solutions** for h_{11} are

$$\begin{aligned} h_{11} = & \frac{Ar(5 - 3x^2 + 13x^4 + 5x^6)}{5m(1 + x^2)^2} - \frac{3r^2(1 - x^2)}{2m^2(1 + x^2)^3} \left[-K_1^{(2)}(1 - x^2) + 2K_3^{(2)}x \right] \\ & + \frac{A^2r^2}{300m^2(1 + x^2)^4} (-675 + 92x + 2752x^2 - 1851x^4 - 92x^5 - 268x^6 + 747x^8 \\ & + 420x^{10} + 75x^{12}) + \mathcal{O}(r^3) + \dots \end{aligned}$$

To locate the **extremal Kerr black hole** in the **moduli space of deformations**, we can either:

- Compare it to the **next-to-lowest order** solutions to fix A , $K_1^{(2)}$ and $K_3^{(2)}$.
- Projects it onto $H^1(V)$ using p_1 , then compare it to the **lowest-order** solution to fix all A , $K_1^{(n)}$ and $K_3^{(n)}$.

We find

$$A = \frac{15}{7}, \quad K_1^{(2)} = \frac{277}{49}, \quad K_3^{(2)} = \frac{23}{49}, \quad K_1^{(3)} = \frac{2684}{1029}, \quad K_3^{(3)} = -\frac{254}{343}.$$

Summary and outlook

- The deformation problem of near-horizon geometry is encoded as an L_∞ -algebra. The Einstein field equations are the Maurer–Cartan equation.
- The problem reduces to solving the Maurer–Cartan equation of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- In extremal Kerr near-horizon geometry, the equation is algebraic and trivially satisfied (up to the next-to-lowest order) for all $\Theta^\circ \in H^1(V)$.
- The formalism gives an upper bound for the degrees of freedom of the moduli space of deformations, which can be lowered by including higher-order terms in the Maurer–Cartan equation of the minimal model.
- Explore warp-product of AdS space can be expressed in Gaussian-null coordinates (connection to holography)
- Investigate non-uniqueness in 5D bubbling solutions [Horowitz, Kunduri, and Lucietti].

Summary and outlook

- The deformation problem of near-horizon geometry is encoded as an L_∞ -algebra. The Einstein field equations are the Maurer–Cartan equation.
- The problem reduces to solving the Maurer–Cartan equation of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- In extremal Kerr near-horizon geometry, the equation is algebraic and trivially satisfied (up to the next-to-lowest order) for all $\Theta^\circ \in H^1(V)$.
- The formalism gives an upper bound for the degrees of freedom of the moduli space of deformations, which can be lowered by including higher-order terms in the Maurer–Cartan equation of the minimal model.
- Explore warp-product of AdS space can be expressed in Gaussian-null coordinates (connection to holography)
- Investigate non-uniqueness in 5D bubbling solutions [Horowitz, Kunduri, and Lucietti].

Summary and outlook

- The deformation problem of near-horizon geometry is encoded as an L_∞ -algebra. The Einstein field equations are the Maurer–Cartan equation.
- The problem reduces to solving the Maurer–Cartan equation of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- In extremal Kerr near-horizon geometry, the equation is algebraic and trivially satisfied (up to the next-to-lowest order) for all $\Theta^\circ \in H^1(V)$.
- The formalism gives an upper bound for the degrees of freedom of the moduli space of deformations, which can be lowered by including higher-order terms in the Maurer–Cartan equation of the minimal model.
- Explore warp-product of AdS space can be expressed in Gaussian-null coordinates (connection to holography)
- Investigate non-uniqueness in 5D bubbling solutions [Horowitz, Kunduri, and Lucietti].

Summary and outlook

- The deformation problem of near-horizon geometry is encoded as an L_∞ -algebra. The Einstein field equations are the Maurer–Cartan equation.
- The problem reduces to solving the Maurer–Cartan equation of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- In extremal Kerr near-horizon geometry, the equation is algebraic and trivially satisfied (up to the next-to-lowest order) for all $\Theta^\circ \in H^1(V)$.
- The formalism gives an upper bound for the degrees of freedom of the moduli space of deformations, which can be lowered by including higher-order terms in the Maurer–Cartan equation of the minimal model.
- Explore warp-product of AdS space can be expressed in Gaussian-null coordinates (connection to holography)
- Investigate non-uniqueness in 5D bubbling solutions [Horowitz, Kunduri, and Lucietti].

Summary and outlook

- The deformation problem of near-horizon geometry is encoded as an L_∞ -algebra. The Einstein field equations are the Maurer–Cartan equation.
- The problem reduces to solving the Maurer–Cartan equation of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- In extremal Kerr near-horizon geometry, the equation is algebraic and trivially satisfied (up to the next-to-lowest order) for all $\Theta^\circ \in H^1(V)$.
- The formalism gives an upper bound for the degrees of freedom of the moduli space of deformations, which can be lowered by including higher-order terms in the Maurer–Cartan equation of the minimal model.
- Explore warp-product of AdS space can be expressed in Gaussian-null coordinates (connection to holography)
- Investigate non-uniqueness in 5D bubbling solutions [Horowitz, Kunduri, and Lucietti].

Summary and outlook

- The deformation problem of near-horizon geometry is encoded as an L_∞ -algebra. The Einstein field equations are the Maurer–Cartan equation.
- The problem reduces to solving the Maurer–Cartan equation of the minimal model $(H^\bullet(V), \mu_i^\circ)$.
- In extremal Kerr near-horizon geometry, the equation is algebraic and trivially satisfied (up to the next-to-lowest order) for all $\Theta^\circ \in H^1(V)$.
- The formalism gives an upper bound for the degrees of freedom of the moduli space of deformations, which can be lowered by including higher-order terms in the Maurer–Cartan equation of the minimal model.
- Explore warp-product of AdS space can be expressed in Gaussian-null coordinates (connection to holography)
- Investigate non-uniqueness in 5D bubbling solutions [Horowitz, Kunduri, and Lucietti].