

Semigroups Of Generalised Symmetries

Yann Péresse

Department of Physics, Astronomy, and Mathematics
University of Hertfordshire

Mathematics and Theoretical Physics Seminar
9th of January 2026



University of
Hertfordshire **UH**



The concept of “symmetry” originally belongs in Geometry.

Example: A rectangle has four symmetries:




id_X



σ_v



σ_h



ρ_π

The symmetries of a geometric shape X are the **distance preserving** bijections $f : X \rightarrow X$.

The symmetries of any object X are the bijections $f : X \rightarrow X$ which preserve the structure of X .



object X	Symmetries of X	preserved structure
metric space	isometries	distance
topological space	homeomorphisms	open sets
relational structure	automorphisms	tuples
set	all bijections	only cardinality

The symmetries of any object X are the bijections $f : X \rightarrow X$ which preserve the structure of X .



object X	Symmetries of X	preserved structure
metric space	isometries	distance
topological space	homeomorphisms	open sets
relational structure	automorphisms	tuples
set	all bijections	only cardinality

The symmetries of any X form a subgroup of the symmetric group $\text{Sym}(X)$.

Three shapes with the same symmetry groups:





Three shapes with the same symmetry groups:





As far as groups are concerned, T , T' , and S are the same.

Three shapes with the same symmetry groups:



As far as groups are concerned, T , T' , and S are the same.
Is this a feature or a bug?



Three shapes with the same symmetry groups:



As far as groups are concerned, T , T' , and S are the same.
Is this a feature or a bug?

- + Maybe, intuitively, T' is not a hexagon but a “triangle with cut-off corners”.



Three shapes with the same symmetry groups:



As far as groups are concerned, T , T' , and S are the same.
Is this a feature or a bug?

- + Maybe, intuitively, T' is not a hexagon but a “triangle with cut-off corners”.
- + And the Sierpiński **triangle** S is very “triangly”.

Three shapes with the same symmetry groups:



As far as groups are concerned, T , T' , and S are the same.
Is this a feature or a bug?

- + Maybe, intuitively, T' is not a hexagon but a “triangle with cut-off corners”.
- + And the Sierpiński **triangle** S is very “triangly”.
- But, also intuitively, S seems to have a lot more symmetry (self-similarity). This is not captured by groups.

Transformations and Partial bijections

Let X be a set. We may extend the symmetric group $\text{Sym}(X)$ in two different ways.

Transformations and Partial bijections

Let X be a set. We may extend the symmetric group $\text{Sym}(X)$ in two different ways.

- A *transformation* of X is a function $f : X \rightarrow X$.
- A *partial bijection* on X is a bijection $f : Y \rightarrow Z$ where Y and Z are subsets of X .

Transformations and Partial bijections

Let X be a set. We may extend the symmetric group $\text{Sym}(X)$ in two different ways.

- A *transformation* of X is a function $f : X \rightarrow X$.
- A *partial bijection* on X is a bijection $f : Y \rightarrow Z$ where Y and Z are subsets of X .
- The *Full Transformation Semigroup* X^X consists of all transformations of X .
- The *Symmetric Inverse Semigroup* I_X consists of all partial bijections on X .

Recall

Recall (or see for the first time):

Some abstract definitions. . .

Recall (or see for the first time):

- ***semigroup***: a set with an associative binary operation.

Some abstract definitions. . .


Recall (or see for the first time):

- ***semigroup***: a set with an associative binary operation.
- ***monoid***: a semigroup with an identity element.

Some abstract definitions...

Recall (or see for the first time):

- **semigroup**: a set with an associative binary operation.
- **monoid**: a semigroup with an identity element.
- **inverse semigroup**: a semigroup S such that $(\forall x \in S)(\exists !x^{-1} \in S) : xx^{-1}x = x$ and $x^{-1}xx^{-1} = x^{-1}$.



... and why you may immediately forget them.

Theorem (A. Cayley, 1854)

Every group is isomorphic to a subgroup of some $\text{Sym}(X)$.

Theorem (Folklore, 20th century)

Every semigroup is isomorphic to a subsemigroup of some X^X .

Theorem (Wagner-Preston, 1950s)

Every inverse semigroup is isomorphic to an inverse subsemigroup of some I_X .

... and why you may immediately forget them.

Theorem (A. Cayley, 1854)

Every group is isomorphic to a subgroup of some $\text{Sym}(X)$.

Theorem (Folklore, 20th century)

Every semigroup is isomorphic to a subsemigroup of some X^X .

Theorem (Wagner-Preston, 1950s)

Every inverse semigroup is isomorphic to an inverse subsemigroup of some I_X .

The rest of this talk is mostly about $\text{Sym}(X)$, X^X , and I_X when X is an infinite set.

- **Serhii Bardyla**, University of Vienna, Austria
- **Igor Dolinka**, University of Novi Sad, Serbia
- **James East**, Western Sydney University, Australia
- **Luna Elliott**, University of Manchester, UK
- **Martin Hampenberg**, University of Hertfordshire, UK
- **James Hyde**, Cornell University, USA
- **Julius Jonušas**, enspired GmbH, Austria
- **Zak Mesyan**, University of Colorado, USA
- **James D. Mitchell**, University of St Andrews, UK
- **Michał Morayne**, Technical University of Wrocław, Poland
- **Michael Pinsker**, Technical University of Vienna, Austria
- **Martin Quick**, University of St Andrews, UK

If X is a finite set, then

$$\text{Sym}(X) = |X|! \quad |X^X| = |X|^{|X|} \quad |I_X| = \sum_{k=0}^{|X|} \binom{|X|}{k}^2 k!$$

If X is a finite set, then

$$\text{Sym}(X) = |X|! \quad |X^X| = |X|^{|X|} \quad |I_X| = \sum_{k=0}^{|X|} \binom{|X|}{k}^2 k!$$

If X is infinite, then

$$|\text{Sym}(X)| = |X^X| = |I_X| = 2^{|X|}.$$

If X is a finite set, then

$$\text{Sym}(X) = |X|! \quad |X^X| = |X|^{|X|} \quad |I_X| = \sum_{k=0}^{|X|} \binom{|X|}{k}^2 k!$$

If X is infinite, then

$$|\text{Sym}(X)| = |X^X| = |I_X| = 2^{|X|}.$$

In the smallest infinite case $X = \mathbb{N} = \{0, 1, 2, 3, \dots\}$,

$$|\text{Sym}(\mathbb{N})| = |\mathbb{N}^{\mathbb{N}}| = |I_{\mathbb{N}}| = 2^{\aleph_0} = \mathfrak{c}$$

where $\mathfrak{c} = |\mathbb{R}|$ is the *continuum*.

The *rank* of a semigroup S is the least cardinality of a generating set for S .

The *rank* of a semigroup S is the least cardinality of a generating set for S .

If X is finite with $|X| \geq 3$, then

- $\text{Sym}(X)$ has rank 2.
- X^X has rank 3.
- I_X has rank 3.

The *rank* of a semigroup S is the least cardinality of a generating set for S .

If X is finite with $|X| \geq 3$, then

- $\text{Sym}(X)$ has rank 2.
- X^X has rank 3.
- I_X has rank 3.

What about the infinite case?

The *rank* of a semigroup S is the least cardinality of a generating set for S .

If X is finite with $|X| \geq 3$, then

- $\text{Sym}(X)$ has rank 2.
- X^X has rank 3.
- I_X has rank 3.

What about the infinite case?

- If X is infinite, then $\text{Sym}(X)$, X^X , and I_X are uncountable.

The *rank* of a semigroup S is the least cardinality of a generating set for S .

If X is finite with $|X| \geq 3$, then

- $\text{Sym}(X)$ has rank 2.
- X^X has rank 3.
- I_X has rank 3.

What about the infinite case?

- If X is infinite, then $\text{Sym}(X)$, X^X , and I_X are uncountable.
- Every uncountable semigroup S has rank $|S|$.

The *rank* of a semigroup S is the least cardinality of a generating set for S .

If X is finite with $|X| \geq 3$, then

- $\text{Sym}(X)$ has rank 2.
- X^X has rank 3.
- I_X has rank 3.

What about the infinite case?

- If X is infinite, then $\text{Sym}(X)$, X^X , and I_X are uncountable.
- Every uncountable semigroup S has rank $|S|$.

Is there a better notion of rank for uncountable semigroups?

The *Sierpiński rank* of a semigroup S is the least $n \in \mathbb{N} \cup \{\infty\}$ such that every countable subset of S is contained in an n -generated subsemigroup of S .

Sierpiński rank

The *Sierpiński rank* of a semigroup S is the least $n \in \mathbb{N} \cup \{\infty\}$ such that every countable subset of S is contained in an n -generated subsemigroup of S .

If X is any infinite set, then:

Theorem (W. Sierpiński, 1935)

The Sierpiński rank of X^X is 2.

Theorem (F. Galvin, 1995)

The Sierpiński rank of $\text{Sym}(X)$ is 2.

Theorem (J. Hyde, YP, 2012)

The Sierpiński rank of I_X is 2.

Other semigroups with finite Sierpiński rank

- rank 2 The **homeomorphism groups** of certain **totally disconnected spaces** including the rational numbers, the Baire space, and the Cantor space. (F. Galvin's proof)
- rank 2 the **continuous functions** on any of the **rational numbers** \mathbb{Q} , the m -dimensional **unit square** $[0, 1]^m$, or the **Hilbert cube** $[0, 1]^{\mathbb{N}}$ (Cook, Ingram, 1969);
- rank 2 the **linear maps** on an **infinite dimensional vector space** (Magill, 1988);
- rank 3 the **order endomorphisms** of the **unit interval** $[0, 1]$ (Mitchell, YP, Quick, 2007).
- rank 2 the **endomorphisms** of certain **Fraïssé limits** such as the **Random Graph** (Dolinka, YP, 2014);
- rank 2 The **order automorphisms** of the rational numbers \mathbb{Q} (J. Hyde, J. Jonusas, J.D. Mitchell, YP, 2016)

Consequences of finite Sierpiński rank 1: embeddings

From the facts that $\text{Sym}(X)$, X^X , and $I_X \dots$

- 1 ... have Sierpiński rank 2;
- 2 ... contain all groups, semigroups, and inverse semigroups, respectively;

we get the following corollary.

Corollary

Every countable group (semigroup, inverse semigroup) is a subgroup (subsemigroup, inverse subsemigroup) of a 2-generator group (semigroup, inverse semigroup).

Definition (relative rank)

Let T be a subsemigroup of S . The *relative rank* of T in S is the least cardinality of a subset A of S such that $A \cup T$ generates S .

Definition (relative rank)

Let T be a subsemigroup of S . The *relative rank* of T in S is the least cardinality of a subset A of S such that $A \cup T$ generates S .

Relative rank is an algebraic measure of the size of a subsemigroup T . Big subsemigroups have small relative rank.

Definition (relative rank)

Let T be a subsemigroup of S . The *relative rank* of T in S is the least cardinality of a subset A of S such that $A \cup T$ generates S .

Relative rank is an algebraic measure of the size of a subsemigroup T . Big subsemigroups have small relative rank.

Proposition

Let S be a semigroup with finite Sierpiński rank n and T a subsemigroup of S . Then the relative rank of T in S is either at most n or uncountable.

Subgroups and subsemigroups

Subgroups and subsemigroups

Let X be infinite. Then studying substructures of $\text{Sym}(X)$, X^X , and I_X is hard:

Subgroups and subsemigroups

Let X be infinite. Then studying substructures of $\text{Sym}(X)$, X^X , and I_X is hard:

Theorem (R. Ball, 1966)

$\text{Sym}(X)$ has $2^{(2^{|X|})}$ many maximal subgroups.

Theorem (J. East, J.D. Mitchell, YP, 2011)

X^X has $2^{(2^{|X|})}$ many maximal subsemigroups.

Theorem (M. Hampenberg, YP, 2025)

I_X has $2^{(2^{|X|})}$ many maximal inverse subsemigroups.

Subgroups and subsemigroups

Let X be infinite. Then studying substructures of $\text{Sym}(X)$, X^X , and I_X is hard:

Theorem (R. Ball, 1966)

$\text{Sym}(X)$ has $2^{(2^{|X|})}$ many maximal subgroups.

Theorem (J. East, J.D. Mitchell, YP, 2011)

X^X has $2^{(2^{|X|})}$ many maximal subsemigroups.

Theorem (M. Hampenberg, YP, 2025)

I_X has $2^{(2^{|X|})}$ many maximal inverse subsemigroups.

So $\text{Sym}(X)$, I_X , and X^X have as many maximal substructures as they have subsets!

Subgroups and subsemigroups

Let X be infinite. Then studying substructures of $\text{Sym}(X)$, X^X , and I_X is hard:

Theorem (R. Ball, 1966)

$\text{Sym}(X)$ has $2^{(2^{|X|})}$ many maximal subgroups.

Theorem (J. East, J.D. Mitchell, YP, 2011)

X^X has $2^{(2^{|X|})}$ many maximal subsemigroups.

Theorem (M. Hampenberg, YP, 2025)

I_X has $2^{(2^{|X|})}$ many maximal inverse subsemigroups.

So $\text{Sym}(X)$, I_X , and X^X have as many maximal substructures as they have subsets!

Let us borrow some extra tools from Topology.

- A *semigroup topology* for a semigroup (S, \cdot) is any topology on S under which the multiplication map

$$(a, b) \mapsto a \cdot b$$

is continuous.

- A *semigroup topology* for a semigroup (S, \cdot) is any topology on S under which the multiplication map

$$(a, b) \mapsto a \cdot b$$

is continuous.

- An *inverse semigroup topology* for an inverse semigroup (I, \cdot) is any topology on I under which the maps

$$(a, b) \mapsto a \cdot b \quad \text{and} \quad a \mapsto a^{-1}$$

are continuous.

- An inverse semigroup topology on a group is called a *group topology*.

Why Topological Algebra?

Some good ways of using Topology in Semigroup Theory:

- 1 Fix a semigroup S . What kind of topologies does S admit?
- 2 Conversely, fix some topological properties (say compact & Hausdorff). What can you say about semigroups admitting such topologies?

Why Topological Algebra?

Some good ways of using Topology in Semigroup Theory:

- 1 Fix a semigroup S . What kind of topologies does S admit?
- 2 Conversely, fix some topological properties (say compact & Hausdorff). What can you say about semigroups admitting such topologies?
- 3 Fix a semigroup S and a semigroup topology τ for S . Study topologically-algebraic problems:
 - What are the subsemigroups of S which are closed (or open, compact, ...) under τ ?
 - What is the least number of elements of S that generates a subsemigroup of S which is dense under τ ? (“topologically generating S ”)

Why Topological Algebra?

Some good ways of using Topology in Semigroup Theory:

- 1 Fix a semigroup S . What kind of topologies does S admit?
- 2 Conversely, fix some topological properties (say compact & Hausdorff). What can you say about semigroups admitting such topologies?
- 3 Fix a semigroup S and a semigroup topology τ for S . Study topologically-algebraic problems:
 - What are the subsemigroups of S which are closed (or open, compact, ...) under τ ?
 - What is the least number of elements of S that generates a subsemigroup of S which is dense under τ ? (“topologically generating S ”)

For point 3 to be interesting and meaningful, we need to agree on a τ which is (i) ‘nice’ in a topological sense and (ii) ‘natural’ for S .

Properties that make topologists happy

Part 1. Having **enough** open sets

Properties that make topologists happy

Part 1. Having **enough** open sets

There are nine, increasingly stronger, “separation axioms”

$$T_0 \Leftarrow T_1 \Leftarrow T_2 \Leftarrow T_{2\frac{1}{2}} \Leftarrow T_3 \Leftarrow T_{3\frac{1}{2}} \Leftarrow T_4 \Leftarrow T_5 \Leftarrow T_6.$$

They describe ways in which points in the space may be separated by open sets.

Properties that make topologists happy

Part 1. Having **enough** open sets

There are nine, increasingly stronger, “separation axioms”

$$T_0 \Leftarrow T_1 \Leftarrow T_2 \Leftarrow T_{2\frac{1}{2}} \Leftarrow T_3 \Leftarrow T_{3\frac{1}{2}} \Leftarrow T_4 \Leftarrow T_5 \Leftarrow T_6.$$

They describe ways in which points in the space may be separated by open sets. For example, a topological space S is...

- ... T_1 (Fréchet) if for all distinct $x, y \in S$, there exists an open neighbourhood of x which does not contain y ;
- ... T_2 (Hausdorff) if all distinct $x, y \in S$ have disjoint open neighbourhoods.

$U \subseteq S$ is a *neighbourhood* of $x \in S$ if $x \in V \subseteq U$ for some open $V \subseteq S$.

Properties that make topologists happy

Part 2. Having **not too many** open sets

Properties that make topologists happy

Part 2. Having **not too many** open sets

A topological space S is...

- ... *separable* if S has a countable dense subset;

Properties that make topologists happy

Part 2. Having **not too many** open sets

A topological space S is...

- ... *separable* if S has a countable dense subset;
- ... *compact* if every open cover of S may be reduced to a finite subcover;
- ... *connected* if no open set (other than \emptyset and S) is also closed;

Properties that make topologists happy

Part 3. Being like \mathbb{R}

Properties that make topologists happy

Part 3. Being like \mathbb{R}

A topological space (S, τ) is...

- ... *second-countable* if τ has a countable basis;
- ... *metrizable* if τ is induced by a metric on S ;
- ... *completely metrizable* if τ is induced by a complete metric on S ;
- ... *Polish* if S is completely metrizable and separable.
- ... *locally compact* if every point in S has a compact neighbourhood.

Natural topologies for a semigroup

(S, \cdot) is a semigroup. What semigroup topology should we give S ?

Two approaches:

Natural topologies for a semigroup

(S, \cdot) is a semigroup. What semigroup topology should we give S ?

Two approaches:

- 1 **From context:** What kind of object is S ? Does the set S already come with a topology we care about?

Natural topologies for a semigroup

(S, \cdot) is a semigroup. What semigroup topology should we give S ?

Two approaches:

- 1 **From context:** What kind of object is S ? Does the set S already come with a topology we care about? Examples:
 - the real numbers under addition $(\mathbb{R}, +)$
 - general linear groups $GL_n(\mathbb{R})$

Natural topologies for a semigroup

(S, \cdot) is a semigroup. What semigroup topology should we give S ?

Two approaches:

- 1 **From context:** What kind of object is S ? Does the set S already come with a topology we care about? Examples:
 - the real numbers under addition $(\mathbb{R}, +)$
 - general linear groups $GL_n(\mathbb{R})$
- 2 **Purely algebraic:** Ignore the context (if any) of the set S as an object and consider topologies that may be defined on any abstract semigroup (S, \cdot) .

Natural topologies for a semigroup

(S, \cdot) is a semigroup. What semigroup topology should we give S ?

Two approaches:

- 1 **From context:** What kind of object is S ? Does the set S already come with a topology we care about? Examples:
 - the real numbers under addition $(\mathbb{R}, +)$
 - general linear groups $GL_n(\mathbb{R})$
- 2 **Purely algebraic:** Ignore the context (if any) of the set S as an object and consider topologies that may be defined on any abstract semigroup (S, \cdot) . Examples:
 - Minimal topologies which are T_1 , Hausdorff, . . .
 - Maximal topologies which are compact, second-countable, . . .
 - Topologies defined via algebraic equations (Zariski topologies).

A topology for $\mathbb{N}^{\mathbb{N}}$ and $\text{Sym}(\mathbb{N})$

The *pointwise topology* is canonical:

A topology for $\mathbb{N}^{\mathbb{N}}$ and $\text{Sym}(\mathbb{N})$

The *pointwise topology* is canonical:

- Naturally defined as the product topology on $\mathbb{N}^{\mathbb{N}}$;
- $\text{Sym}(\mathbb{N})$ receives the subspace topology;

A topology for $\mathbb{N}^{\mathbb{N}}$ and $\text{Sym}(\mathbb{N})$

The *pointwise topology* is canonical:

- Naturally defined as the product topology on $\mathbb{N}^{\mathbb{N}}$;
- $\text{Sym}(\mathbb{N})$ receives the subspace topology;
- Coincides with the *Zariski topologies* on $\text{Sym}(\mathbb{N})$ and $\mathbb{N}^{\mathbb{N}}$.

A topology for $\mathbb{N}^{\mathbb{N}}$ and $\text{Sym}(\mathbb{N})$

The *pointwise topology* is canonical:

- Naturally defined as the product topology on $\mathbb{N}^{\mathbb{N}}$;
- $\text{Sym}(\mathbb{N})$ receives the subspace topology;
- Coincides with the *Zariski topologies* on $\text{Sym}(\mathbb{N})$ and $\mathbb{N}^{\mathbb{N}}$.
- $\mathbb{N}^{\mathbb{N}}$ is a topological semigroup;
- $\text{Sym}(\mathbb{N})$ is a topological group;

A topology for $\mathbb{N}^{\mathbb{N}}$ and $\text{Sym}(\mathbb{N})$

The *pointwise topology* is canonical:

- Naturally defined as the product topology on $\mathbb{N}^{\mathbb{N}}$;
- $\text{Sym}(\mathbb{N})$ receives the subspace topology;
- Coincides with the *Zariski topologies* on $\text{Sym}(\mathbb{N})$ and $\mathbb{N}^{\mathbb{N}}$.
- $\mathbb{N}^{\mathbb{N}}$ is a topological semigroup;
- $\text{Sym}(\mathbb{N})$ is a topological group;
- $\mathbb{N}^{\mathbb{N}}$ and $\text{Sym}(\mathbb{N})$ are Polish (completely metrizable & separable);

A topology for $\mathbb{N}^{\mathbb{N}}$ and $\text{Sym}(\mathbb{N})$

The *pointwise topology* is canonical:

- Naturally defined as the product topology on $\mathbb{N}^{\mathbb{N}}$;
- $\text{Sym}(\mathbb{N})$ receives the subspace topology;
- Coincides with the *Zariski topologies* on $\text{Sym}(\mathbb{N})$ and $\mathbb{N}^{\mathbb{N}}$.
- $\mathbb{N}^{\mathbb{N}}$ is a topological semigroup;
- $\text{Sym}(\mathbb{N})$ is a topological group;
- $\mathbb{N}^{\mathbb{N}}$ and $\text{Sym}(\mathbb{N})$ are Polish (completely metrizable & separable);
- $M \leq \mathbb{N}^{\mathbb{N}}$ is closed $\iff M = \text{End}(R)$ for some relational structure R on \mathbb{N} ;
- $G \leq \text{Sym}(\mathbb{N})$ is closed $\iff G = \text{Aut}(R)$ for some relational structure R on \mathbb{N} .

Unique topologies: $\text{Sym}(\mathbb{N})$

Theorem (Gaughan 1967)

The pointwise topology is the least Hausdorff group topology on $\text{Sym}(\mathbb{N})$.

Theorem (Kechris, Rosendal 2004)

- *The pointwise topology is the unique non-trivial separable group topology on $\text{Sym}(\mathbb{N})$.*
- *Every homomorphism from $\text{Sym}(\mathbb{N})$ to a separable group is continuous (automatic continuity).*

Corollary

The pointwise topology is the unique Polish group topology on $\text{Sym}(\mathbb{N})$.

Unique topologies: $\text{Sym}(\mathbb{N})$

Theorem (Gaughan 1967)

The pointwise topology is the least Hausdorff group topology on $\text{Sym}(\mathbb{N})$.

Theorem (Kechris, Rosendal 2004)

- *The pointwise topology is the unique non-trivial separable group topology on $\text{Sym}(\mathbb{N})$.*
- *Every homomorphism from $\text{Sym}(\mathbb{N})$ to a separable group is continuous (automatic continuity).*

Corollary

The pointwise topology is the unique Polish group topology on $\text{Sym}(\mathbb{N})$.

What about $\mathbb{N}^{\mathbb{N}}$ and $I_{\mathbb{N}}$?

Unique topologies: $\mathbb{N}^{\mathbb{N}}$ and friends

Theorem (L. Elliott, J. Jonušas, J.D. Mitchell, Z. Mesyan, M. Morayne, YP 2023)

The following transformation monoids have a unique Polish semigroup topology:

- $\mathbb{N}^{\mathbb{N}}$;
- *The monoid $P(\mathbb{N})$ of partial transformations of \mathbb{N} ;*
- *The monoid $C([0, 1]^{\mathbb{N}})$ of continuous transformations of the Hilbert cube $[0, 1]^{\mathbb{N}}$;*
- *The monoid $C(2^{\mathbb{N}})$ of continuous transformations of the Cantor set $2^{\mathbb{N}}$.*

Moreover, $\mathbb{N}^{\mathbb{N}}$, $P(\mathbb{N})$, and $C(2^{\mathbb{N}})$ have automatic continuity: every homomorphism to a second-countable semigroup is continuous.

Unique topologies: endomorphism semigroups

Theorem (L. Elliott, J. Jonušas, J.D. Mitchell, YP, M. Pinsker 2023)

The pointwise topology is the unique Polish semigroup topology on the endomorphism monoids of the following relational structures:

- 1 *the random graph;*
- 2 *the random directed graph;*
- 3 *infinitely many copies of a finite complete graph K_n ;*
- 4 *a partition of \mathbb{N} into finitely many infinite classes;*
- 5 *the random strict partial order.*

The endomorphism monoids of 1-4 have automatic continuity.

Unique (?) topologies: $I_{\mathbb{N}}$

Theorem (S. Bardyla, L. Elliott, J. Jonušas, J.D. Mitchell, Z. Mesyan, M. Morayne, YP 2023)

The symmetric inverse semigroup $I_{\mathbb{N}}$ has

Unique (?) topologies: $I_{\mathbb{N}}$

Theorem (S. Bardyla, L. Elliott, J. Jonušas, J.D. Mitchell, Z. Mesyan, M. Morayne, YP 2023)

The symmetric inverse semigroup $I_{\mathbb{N}}$ has

- *countably infinitely many Polish semigroup topologies (which can be classified);*

Theorem (S. Bardyla, L. Elliott, J. Jonušas, J.D. Mitchell, Z. Mesyan, M. Morayne, YP 2023)

The symmetric inverse semigroup $I_{\mathbb{N}}$ has

- *countably infinitely many Polish semigroup topologies (which can be classified);*
- *two minimal Polish semigroup topologies;*

Unique (?) topologies: $I_{\mathbb{N}}$

Theorem (S. Bardyla, L. Elliott, J. Jonušas, J.D. Mitchell, Z. Mesyan, M. Morayne, YP 2023)

The symmetric inverse semigroup $I_{\mathbb{N}}$ has

- *countably infinitely many Polish semigroup topologies (which can be classified);*
- *two minimal Polish semigroup topologies;*
- *one maximal Polish semigroup topology;*

Unique (?) topologies: $I_{\mathbb{N}}$

Theorem (S. Bardyla, L. Elliott, J. Jonušas, J.D. Mitchell, Z. Mesyan, M. Morayne, YP 2023)

The symmetric inverse semigroup $I_{\mathbb{N}}$ has

- *countably infinitely many Polish semigroup topologies (which can be classified);*
- *two minimal Polish semigroup topologies;*
- *one maximal Polish semigroup topology;*
- *a unique Polish inverse semigroup topology (the maximal Polish semigroup topology).*

Every finite partial order embeds into the partial order of Polish semigroup topologies on $I_{\mathbb{N}}$.

Do closed subsemigroups of $I_{\mathbb{N}}$ correspond to relational structures like they do in $\text{Sym}(\mathbb{N})$ and $\mathbb{N}^{\mathbb{N}}$?

Do closed subsemigroups of $I_{\mathbb{N}}$ correspond to relational structures like they do in $\text{Sym}(\mathbb{N})$ and $\mathbb{N}^{\mathbb{N}}$?

The *partial automorphism monoid* of a relational structure R on \mathbb{N} consists of all isomorphisms between substructures of R .

Do closed subsemigroups of $I_{\mathbb{N}}$ correspond to relational structures like they do in $\text{Sym}(\mathbb{N})$ and $\mathbb{N}^{\mathbb{N}}$?

The *partial automorphism monoid* of a relational structure R on \mathbb{N} consists of all isomorphisms between substructures of R . Every partial automorphism monoid must be *full*; that is closed under taking restrictions of functions.

Do closed subsemigroups of $I_{\mathbb{N}}$ correspond to relational structures like they do in $\text{Sym}(\mathbb{N})$ and $\mathbb{N}^{\mathbb{N}}$?

The *partial automorphism monoid* of a relational structure R on \mathbb{N} consists of all isomorphisms between substructures of R . Every partial automorphism monoid must be *full*; that is closed under taking restrictions of functions.

Theorem (M. Hampenberg, YP 2024)

Let M be a full inverse submonoid of $I_{\mathbb{N}}$. Then the following are equivalent:

- *M is closed in a Polish semigroup topology on $I_{\mathbb{N}}$;*
- *M is closed in every Polish semigroup topology on $I_{\mathbb{N}}$;*
- *M is the partial automorphism monoid of a relational structure defined on \mathbb{N} .*

The Bergman-Shelah preorder

Recall: The relative rank of a subsemigroup T of S is the least number of extra elements needed to turn T into a generating set for S .

The Bergman-Shelah preorder

Recall: The relative rank of a subsemigroup T of S is the least number of extra elements needed to turn T into a generating set for S .

Theorem (F. Galvin, 1995)

Let X be an infinite set. Then the relative rank of any proper subgroup of $\text{Sym}(X)$ is either 1 or uncountable.

The Bergman-Shelah preorder

Recall: The relative rank of a subsemigroup T of S is the least number of extra elements needed to turn T into a generating set for S .

Theorem (F. Galvin, 1995)

Let X be an infinite set. Then the relative rank of any proper subgroup of $\text{Sym}(X)$ is either 1 or uncountable.

Inspired by this dichotomy, Bergman and Shelah introduced the following pre-order on subgroups of $\text{Sym}(X)$:

$$G \preceq H \iff G \subseteq \langle A, H \rangle \text{ for some finite } A \subseteq \text{Sym}(X).$$

The Bergman-Shelah preorder

Recall: The relative rank of a subsemigroup T of S is the least number of extra elements needed to turn T into a generating set for S .

Theorem (F. Galvin, 1995)

Let X be an infinite set. Then the relative rank of any proper subgroup of $\text{Sym}(X)$ is either 1 or uncountable.

Inspired by this dichotomy, Bergman and Shelah introduced the following pre-order on subgroups of $\text{Sym}(X)$:

$$G \preceq H \iff G \subseteq \langle A, H \rangle \text{ for some finite } A \subseteq \text{Sym}(X).$$

Notation:

$$\begin{aligned} G \approx H &\iff G \preceq H \text{ and } H \preceq G \\ G \prec H &\iff G \preceq H \text{ and } H \not\preceq G. \end{aligned}$$

Four classes of closed subgroups of $\text{Sym}(\mathbb{N})$

Theorem (G. Bergman, S. Shelah, 2006)

Every topologically closed subgroup of $\text{Sym}(\mathbb{N})$ is \approx -equivalent to one of the following:

- 1 the entire symmetric group $\text{Sym}(\mathbb{N})$
- 2 the infinite direct product $G_\omega := S_2 \times S_3 \times S_4 \times \dots$ in its natural action on the partition $\{0, 1\}, \{2, 3, 4\}, \{4, 5, 6, 7\}, \dots$
- 3 the infinite direct product $G_2 := S_2 \times S_2 \times S_2 \times \dots$ in its natural action on the partition $\{0, 1\}, \{2, 3\}, \{4, 5\}, \dots$
- 4 the trivial subgroup $\{1_{\mathbb{N}}\}$.

Moreover, $\{1_{\mathbb{N}}\} \prec G_2 \prec G_\omega \prec \text{Sym}(\mathbb{N})$

Relative ranks of closed subgroups of $\text{Sym}(\mathbb{N})$

Using the result of Bergman and Shelah, we can classify the relative ranks of closed subgroups of $\text{Sym}(\mathbb{N})$.

Theorem (J.D. Mitchell, M. Morayne, YP, 2010)

Let G be a proper subgroup of $\text{Sym}(\mathbb{N})$ which is closed in the pointwise topology. Then the relative rank of G in $\text{Sym}(\mathbb{N})$ is one of the following: 1, the dominating number \mathfrak{d} , or the continuum 2^{\aleph_0} .

Theorem (M. Hampenberg, YP, 2024)

Let M be the monoid of automorphisms between substructures of a relational structure R defined on \mathbb{N} . If R has finitely many relations, then $M \approx I_{\mathbb{N}}$.

Theorem (M. Hampenberg, YP, 2024)

Let M be the monoid of automorphisms between substructures of a relational structure R defined on \mathbb{N} . If R has finitely many relations, then $M \approx I_{\mathbb{N}}$.

Further results:

- The Bergman-Shelah preorder on $I_{\mathbb{N}}$ has at least 4 classes containing closed full inverse monoids.
- We conjecture that these are the only ones.

The Bergman-Shelah preorder on $\mathbb{N}^{\mathbb{N}}$

Theorem (Z. Mesyan, J.D. Mitchell, M.Morayne, YP, 2012)

The Bergman-Shelah preorder on $\mathbb{N}^{\mathbb{N}}$ has

The Bergman-Shelah preorder on $\mathbb{N}^{\mathbb{N}}$

Theorem (Z. Mesyan, J.D. Mitchell, M.Morayne, YP, 2012)

The Bergman-Shelah preorder on $\mathbb{N}^{\mathbb{N}}$ has

- *at least \aleph_1 many \approx -classes;*

The Bergman-Shelah preorder on $\mathbb{N}^{\mathbb{N}}$

Theorem (Z. Mesyan, J.D. Mitchell, M.Morayne, YP, 2012)

The Bergman-Shelah preorder on $\mathbb{N}^{\mathbb{N}}$ has

- *at least \aleph_1 many \approx -classes;*
- *at least \aleph_0 classes containing closed submonoids of $\mathbb{N}^{\mathbb{N}}$;*

Theorem (Z. Mesyan, J.D. Mitchell, M.Morayne, YP, 2012)

The Bergman-Shelah preorder on $\mathbb{N}^{\mathbb{N}}$ has

- *at least \aleph_1 many \approx -classes;*
- *at least \aleph_0 classes containing closed submonoids of $\mathbb{N}^{\mathbb{N}}$;*
- *arbitrarily large finite anti-chains of \approx -classes containing closed submonoids of $\mathbb{N}^{\mathbb{N}}$.*

Theorem (Z. Mesyan, J.D. Mitchell, M.Morayne, YP, 2012)

The Bergman-Shelah preorder on $\mathbb{N}^{\mathbb{N}}$ has

- *at least \aleph_1 many \approx -classes;*
- *at least \aleph_0 classes containing closed submonoids of $\mathbb{N}^{\mathbb{N}}$;*
- *arbitrarily large finite anti-chains of \approx -classes containing closed submonoids of $\mathbb{N}^{\mathbb{N}}$.*

Moreover, the following are equivalent.

- 1 *There exists a subsemigroup S of $\mathbb{N}^{\mathbb{N}}$ such that $S \approx \mathbb{N}^{\mathbb{N}}$ and every subsemigroup T of S satisfies $T \approx S$ or $T \approx \{1_{\mathbb{N}}\}$.*
- 2 *The continuum hypothesis.*

Thank you for listening!